

Contractions on a manifold polarized by an ample vector bundle

M. Andreatta - M. Mella

Dipartimento di Matematica, Università di Trento,
38050 Povo (TN), Italia

e-mail : Andreatta or Mella @itnvax.science.unitn.it

February 5, 2008

MSC numbers: 14C20, 14E30, 14J40, 14J45

Introduction

An algebraic variety X of dimension n (over the complex field) together with an ample vector bundle E on it will be called a **generalized polarized variety**. The adjoint bundle of the pair (X, E) is the line bundle $K_X + \det(E)$. Problems concerning adjoint bundles have drawn a lot of attention to algebraic geometry: the classical case is when E is a (direct sum of) line bundle (polarized variety), while the generalized case was motivated by the solution of Hartshorne-Frankel conjecture by Mori ([Mo]) and by consequent conjectures of Mukai ([Mu]).

A first point of view is to study the positivity (the nefness or ampleness) of the adjoint line bundle in the case $r = \text{rank}(E)$ is about $n = \dim X$. This was done in a sequel of papers for $r \geq (n - 1)$ and for smooth manifold X ([Ye-Zhang], [Fujita], [Andreatta-Ballico-Wisniewski]). In this paper we want to discuss the next case, namely when $\text{rank}(E) = (n - 2)$, with X smooth; we obtain a complete answer which is described in the theorem (4.1). This is divided in three cases, namely when $K_X + \det(E)$ is not nef, when it is nef and not big and finally when it is nef and big but not ample. If $n = 3$ a complete picture is already contained in the famous paper of Mori ([Mo1]), while the particular case in which $E = \oplus^{(n-2)}(L)$ with L a line bundle was also studied ([Fu1], [So]; in the singular case see [An]). The part 1 of the theorem was proved (in a slightly weaker form) by Zhang ([Zh]) and, in the case E is spanned by global sections, by Wisniewski ([Wi2]).

Another point of view can be the following: let (X, E) be a generalized polarized variety with X smooth and $\text{rank} E = r$. If $K_X + \det(E)$ is nef, then by the Kawamata-Shokurov base point free theorem it supports a contraction (see (1.2)); i.e. there exists a map $\pi : X \rightarrow W$ from X onto a normal projective variety W with connected fiber and such that $K_X + \det(E) = \pi^* H$ for some ample line bundle H on W . It is not difficult to see that, for every fiber F of π , we have $\dim F \geq (r - 1)$, equality holds only if $\dim X > \dim W$. In the paper we study the "border" cases: we assume that $\dim F = (r - 1)$ for every fibers and we prove that X has a $\mathbf{P}^r$ -bundle structure given by π (theorem (3.2)). We consider also the case in which $\dim F = r$ for every fibers and π is birational, proving that W is smooth and that π is a blow-up of a smooth subvariety (theorem (3.1)). This point of view was discussed in the case $E = \oplus^r L$ in the paper [A-W].

Finally in the section (4) we extend the theorem (3.2) to the singular case, namely for projective variety X with log-terminal singularities. In particular this gives the Mukai's conjecture¹ for singular varieties.

1 Notations and generalities

(1.1) We use the standard notations from algebraic geometry. Our language is compatible with that of [K-M-M] to which we refer constantly. We just explain some special definitions and propositions used frequently.

In particular in this paper X will always stand for a smooth complex projective variety of dimension n . Let $\text{Div}(X)$ the group of Cartier divisors on X ; denote by K_X the canonical divisor of X , an element of $\text{Div}(X)$ such that $\mathcal{O}_X(K_X) = \Omega_X^n$. Let $N_1(X) = \frac{\{1\text{-cycles}\}}{\equiv} \otimes \mathbf{R}$, $N^1(X) = \frac{\{\text{divisors}\}}{\equiv} \otimes \mathbf{R}$ and $\overline{< NE(X) >} = \overline{\{\text{effective 1-cycles}\}}$; the last is a closed cone in $N_1(X)$. Let also $\rho(X) = \dim_{\mathbf{R}} N^1(X) < \infty$.

Suppose that K_X is not nef, that is there exists an effective curve C such that $K_X \cdot C < 0$.

Theorem 1.1 [KMM] *Let X as above and H a nef Cartier divisor such that $F := H^\perp \cap \overline{< NE(X) >} \setminus \{0\}$ is entirely contained in the set $\{Z \in N_1(X) : K_X \cdot Z < 0\}$, where $H^\perp = \{Z : H \cdot Z = 0\}$. Then there exists a projective morphism $\varphi : X \rightarrow W$ from X onto a normal variety W with the following properties:*

- i) For an irreducible curve C in X , $\varphi(C)$ is a point if and only if $H \cdot C = 0$, if and only if $\text{cl}(C) \in F$.*

- ii) φ has only connected fibers
- iii) $H = \varphi^*(A)$ for some ample divisor A on W .
- iv) The image $\varphi^* : \text{Pic}(W) \rightarrow \text{Pic}(X)$ coincides with $\{D \in \text{Pic}(X) : D \cdot C = 0 \text{ for all } C \in F\}$.

Definition 1.2 The following terminology is mostly used ([KMM], definition 3-2-3). Referring to the above theorem,

the map φ is called a **contraction** (or an **extremal contraction**); the set F is an **extremal face**, while the Cartier divisor H is a **supporting divisor** for the map φ (or the face F). If $\dim_{\mathbf{R}} F = 1$ the face F is called an **extremal ray**, while φ is called an **elementary contraction**.

Remark We have also ([Mo1]) that if X has an extremal ray R then there exists a rational curve C on X such that $0 < -K_X \cdot C \leq n+1$ and $R = R[C] := \{D \in NE(X) : D \equiv \lambda C, \lambda \in \mathbf{R}^+\}$. Such a curve is called an **extremal curve**.

Remark Let $\pi : X \rightarrow V$ denote a contraction of an extremal face F , supported by $H = \pi^*A$ ([iii]1.1). Let R be an extremal ray in F and $\rho : X \rightarrow W$ the contraction of R . Since $\pi^*A \cdot R = 0$, π^*A comes from $\text{Pic}(W)$ ([iv]1.1). Thus π factors through ρ .

Definition 1.3 To an extremal ray R we can associate:

- i) its length $l(R) := \min\{-K_X \cdot C; \text{ for } C \text{ rational curve and } C \in R\}$
- ii) the locus $E(R) := \{\text{the locus of the curves whose numerical classes are in } R\} \subset X$.

Definition 1.4 It is usual to divide the elementary contractions associated to an extremal ray R in three types according to the dimension of $E(R)$: more precisely we say that φ is of **fiber type**, respectively **divisorial type**, resp. **flipping type**, if $\dim E(R) = n$, resp. $n-1$, resp. $< n-1$. Moreover an extremal ray is said **not nef** if there exists an effective $D \in \text{Div}(X)$ such that $D \cdot C < 0$.

The following very useful inequality was proved in [Io] and [Wi3].

Proposition 1.5 Let φ the contraction of an extremal ray R , $E'(R)$ be any irreducible component of the exceptional locus and d the dimension of a fiber of the contraction restricted to $E'(R)$. Then

$$\dim E'(R) + d \geq n + l(R) - 1.$$

(1.2) Actually it is very useful to understand when a contraction is elementary or in other words when the locus of two distinct extremal rays are disjoint. For this we will use in this paper the following results.

Proposition 1.6 [BS, Corollary 0.6.1] *Let R_1 and R_2 two distinct not nef extremal rays such that $l(R_1) + l(R_2) > n$. Then $E(R_1)$ and $E(R_2)$ are disjoint.*

Something can be said also if $l(R_1) + l(R_2) = n$:

Proposition 1.7 [Fu3, Theorem 2.4] *Let $\pi : X \rightarrow V$ as above and suppose $n \geq 4$ and $l(R_i) \geq n - 2$. Then the exceptional loci corresponding to different extremal rays, are disjoint with each other.*

Proposition 1.8 [ABW1] *Let $\pi : X \rightarrow W$ be a contraction of a face such that $\dim X > \dim W$. Suppose that for every rational curve C in a general fiber of π we have $-K_X \cdot C \geq (n + 1)/2$. Then π is an elementary contraction except if*

- a) $-K_X \cdot C = (n + 2)/2$ for some rational curve C on X , W is a point, X is a Fano manifold of pseudoindex $(n + 2)/2$ and $\rho(X) = 2$
- b) $-K_X \cdot C = (n + 1)/2$ for some rational curve C , and $\dim W \leq 1$

The following definition is used in the theorem:

Definition 1.9 *Let L be an ample line bundle on X . The pair (X, L) is called a scroll (respectively a quadric fibration, respectively a del Pezzo fibration) over a normal variety Y of dimension m if there exists a surjective morphism with connected fibers $\phi : X \rightarrow Y$ such that*

$$K_X + (n - m + 1)L \approx p^* \mathcal{L}$$

(respectively $K_X + (n - m)L \approx p^ \mathcal{L}$; respectively $K_X + (n - m - 1)L \approx p^* \mathcal{L}$) for some ample line bundle $\mathcal{L}$ on Y . X is called a classical scroll (respectively quadric bundle) over a projective variety Y of dimension r if there exists a surjective morphism $\phi : X \rightarrow Y$ such that every fiber is isomorphic to $\mathbf{P}^{n-r}$ (respectively to a quadric in $\mathbf{P}^{(n-r+1)}$) and if there exists a vector bundle E of rank $(n - r + 1)$ (respectively of rank $n - r + 2$) on Y such that $X \simeq \mathbf{P}(E)$ (respectively exists an embedding of X as a subvariety of $\mathbf{P}(E)$).*

2 A technical construction

Let E be a vector bundle of rank r on X and assume that E is ample, in the sense of Hartshorne.

Remark Let $f : \mathbf{P}^1 \rightarrow X$ be a non constant map, and $C = f(\mathbf{P}^1)$, then $\det E \cdot C \geq r$.

In particular if there exists a curve C such that $(K_X + \det E) \cdot C \leq 0$ (for instance if $(K_X + \det E)$ is not nef) then there exists an extremal ray R such that $l(R) \geq r$.

(2.1) Let $Y = \mathbf{P}(E)$ be the associated projective space bundle, $p : Y \rightarrow X$ the natural map onto X and ξ_E the tautological bundle of Y . Then we have the formula for the canonical bundle $K_Y = p^*(K_X + \det E) - r\xi_E$. Note that p is an elementary contraction; let R be the associated extremal ray.

Assume that $K_X + \det E$ is nef but not ample and that it is the supporting divisor of an elementary contraction $\pi : X \rightarrow W$. Then $\rho(Y/W) = 2$ and $-K_Y$ is $\pi \circ p$ -ample. By the relative Mori theory over W we have that there exists a ray on $NE(Y/W)$, say R_1 , of length $\geq r$, not contracted by p , and a relative elementary contraction $\varphi : Y \rightarrow V$. We have thus the following commutative diagram.

$$\begin{array}{ccc} \mathbf{P}(E) = Y & \xrightarrow{\varphi} & V \\ \downarrow p & & \downarrow \psi \\ X & \xrightarrow{\pi} & W \end{array} \quad (1)$$

where φ and ψ are elementary contractions. Let $w \in W$ and let $F(\pi)_w$ be an irreducible component of $\pi^{-1}(w)$; choose also v in $\psi^{-1}(w)$ and let $F(\varphi)_v$ be an irreducible component of $\varphi^{-1}(v)$ such that $p(F(\varphi)_v) \cap F(\pi)_w \neq \emptyset$; then $p(F(\varphi)_v) \subset F(\pi)_w$. This is true by the commutativity of the diagram. Since p and φ are elementary contractions of different extremal rays we have that $\dim(F(\varphi) \cap F(p)) = 0$, that is curve contracted by φ cannot be contracted by p .

In particular this implies that $\dim p(F(\varphi)_v) = \dim F(\varphi)_v$; therefore

$$\dim F(\varphi)_v \leq \dim F(\pi)_w.$$

Remark If $\dim F(\varphi)_v = \dim F(\pi)_w$, then $\dim F(\psi)_w := \dim(\psi^{-1}(w)) = r - 1$; if this holds for every $w \in W$ then ψ is equidimensional.

Proof. Let Y_w be an irreducible component of $p^{-1}\pi^{-1}(w)$ such that $\varphi(Y_w) = F(\psi)_w$. Then $\dim F(\psi)_w = \dim Y_w - \dim F(\varphi)_v = \dim Y_w - \dim F(\pi)_w = \dim F(p) = (r - 1)$. □

(2.2) Slicing techniques

Let $H = \varphi^*(A)$ be a supporting divisor for φ such that the linear system $|H|$ is base point free. We assume as in (2) that $(K_X + \det E)$ is nef and we refer to the diagram (1). The divisor $K_Y + r\xi_E = p^*(K_X + \det E)$ is nef on Y and therefore $m(K_Y + r\xi_E + aH)$, for $m \gg 0$, $a \in \mathbf{N}$, is also a good supporting divisor for φ . Let Z be a smooth n -fold obtained by intersecting $r - 1$ general divisor from the linear system H , i.e. $Z = H_1 \cap \dots \cap H_{r-1}$ (this is what we call a slicing); let $H_i = \varphi^{-1}A_i$.

Note that the map $\varphi' = \varphi|_Z$ is supported by $m|(K_Y + r\xi_E + a\varphi^*A)|_Z|$, hence, by adjunction, it is supported by $K_Z + rL$, where $L = \xi_{E|Z}$. Let $p' = p|_Z$; by construction p' is finite.

If T is (the normalization of) $\varphi(Z)$ and $\psi' : T \rightarrow W$ is the map obtained restricting ψ then we have from (1) the following diagram

$$\begin{array}{ccc} Z & \xrightarrow{\varphi'} & T \\ \downarrow p' & & \downarrow \psi' \\ X & \xrightarrow{\pi} & W \end{array} \quad (2)$$

In general one has a good comprehension of the map φ' (for instance in the case $r = (n - 2)$ see the results in [Fu1] or in [An]). The goal is to "transfer" the information that we have on φ' to the map π . The following proposition is the major step in this program.

Proposition 2.1 *Assume that ψ is equidimensional (in particular this is the case if for every non trivial fiber we have $\dim F(\varphi) = \dim F(\pi)$). Then W has the same singularities of T .*

Proof. By hypothesis any irreducible reduced component F_i of a non trivial fiber $F(\psi)$ is of dimension $r - 1$; this implies also that $F_i = \varphi(F(p))$ for some fiber of p .

Now, let us follow an argument as in [Fu1, Lemma 2.12]. We can assume that the divisor A is very ample; we will choose $r - 1$ divisors $A_i \in |A|$ as above such that, if $T = \bigcap_i A_i$, then $T \cap \psi^{-1}(w)_{red} = N$ is a reduced 0-cycle and $Z = H_1 \cap \dots \cap H_{r-1}$ is a smooth n -fold, where $H_i = \varphi^{-1}A_i$. This can be done by Bertini theorem. Moreover the number of points in N is given by $A^{r-1} \cdot \psi^{-1}(w)_{red} = \sum_i A^{r-1} \cdot F_i = \sum_i d_i$. Note that, by projection formula, we have $A^{r-1} \cdot F_i = \varphi^* A^{r-1} \cdot F(p)$; moreover, since p is a projective bundle, the last number is constant i.e. $\varphi^* A^{r-1} \cdot F(p) = d$ for all fiber $F(p)$, that is the d_i 's are constant.

Now take a small enough neighborhood U of w , in the metric topology, such that any connected component U_λ of $\psi^{-1}(U) \cap T$ meets $\psi^{-1}(w)$ in a single point. This is possible because $\psi' := \psi|_T : T \rightarrow W$ is proper and finite over w . Let ψ_λ

the restriction of ψ at U_λ and m_λ its degree. Then $\deg \psi' = \sum m_\lambda \geq \sum_i d_i = \sum_i d$ and equality holds if and only if ψ is not ramified at w (remember that $\sum_i d_i$ is the number of U_λ).

The generic $F(\psi)_w$ is irreducible and generically reduced. Note that we can choose $\tilde{w} \in W$ such that $\psi^{-1}(\tilde{w}) = \varphi(F(p))$ and $\deg \psi' = A^{r-1} \cdot \psi^{-1}(\tilde{w})$, the latter is possible by the choice of generic sections of $|A|$. Hence, by projection formula $\deg \psi' = A^{r-1} \cdot \psi^{-1}(\tilde{w}) = \varphi^* A^{r-1} \cdot F(p) = d$, that is $m_\lambda = 1$ and the fibers are irreducible. Since W is normal we can conclude, by Zarisky's Main theorem, that W has the same singularity as T . $\square$

3 Some general applications

As an application of the above construction we will prove the following proposition; the case $r = (n - 1)$ was proved in [ABW2].

Proposition 3.1 *Let X be a smooth projective complex variety and E be an ample vector bundle of rank r on X . Assume that $K_X + \det E$ is nef and big but not ample and let $\pi : X \rightarrow W$ be the contraction supported by $K_X + \det E$. Assume also that π is a divisorial elementary contraction, with exceptional divisor D , and that $\dim F \leq r$ for all fibers F . Then W is smooth, π is the blow up of a smooth subvariety $B := \pi(D)$ and $E = \pi^* E' \otimes [-D]$, for some ample E' on W .*

Proof. Let R be the extremal ray contracted by π and $F := F(\pi)$ a fiber. Then $l(R) \geq r$ and thus $\dim F \geq r$ by proposition (1.5). Hence all the fibers of π have dimension r . Consider the commutative diagram (1); let R_1 be the ray contracted by φ . Since $l(R_1) \geq r$, again by proposition (1.5), we have that $\dim F(\varphi) \geq r$ (note that R_1 is not nef). Therefore, since $\dim F(\varphi) \leq \dim F$, we have that $\dim F(\varphi) = \dim F = r$, $l(R) = l(R_1) = r$ and $\xi_E \cdot C_1 = 1$, where C_1 is a (minimal) curve in the ray R_1 . Via slicing we obtain the map $\varphi' : Z \rightarrow T$ which is supported by $K_Z + r\xi_{E|Z}$. This last map is very well understood: namely by [AW, Th 4.1 (iii)] it follows that T is smooth and φ' is a blow up along a smooth subvariety. By proposition (2.1) also W is smooth. Therefore π is a birational morphism between smooth varieties with exceptional locus a prime divisor and with equidimensional non trivial fibers; by [AW, Corollary 4.11] this implies that π is a blow up of a smooth subvariety in W .

We want to show that $E = \pi^* E' \otimes [-D]$. Let D_1 be the exceptional divisor of φ ; first we claim that $\xi_E + D_1$ is a good supporting divisor for φ . To see this observe that $(\xi_E + D_1) \cdot C_1 = 0$, while $(\xi_E + D_1) \cdot C > 0$ for any curve C with $\varphi(C) \neq pt$ (in fact ξ_E is ample and $D_1 \cdot C \geq 0$ for such a curve). Thus

$\xi_E + D_1 = \varphi^* A$ for some ample $A \in \text{Pic}(V)$; moreover by projection formula $A \cdot l = 1$, for any line l in the fiber of ψ . Hence by Grauert theorem $V = \mathbf{P}(E')$ for some ample vector bundle E' on W . This yields, by the commutativity of diagram (1), to $E \otimes D = p_*(\xi_E + D_1) = p_*\varphi^* A = \pi^*\psi_* A = \pi^* E'$. $\square$

We now want to give a similar proposition for the fiber type case.

Theorem 3.2 *Let X be a smooth projective complex variety and E be an ample vector bundle of rank r on X . Assume that $K_X + \det E$ is nef and let $\pi : X \rightarrow W$ be the contraction supported by $K_X + \det E$. Assume that $r \geq (n+1)/2$ and $\dim F \leq r-1$ for any fiber F of π . Then W is smooth, for any fiber $F \simeq \mathbf{P}^{r-1}$ and $E|_F = \oplus^r \mathcal{O}(1)$.*

Proof. Note that by proposition (1.5) π is a contraction of fiber type and all the fibers have dimension $r-1$. Moreover the contraction is elementary, as it follows from proposition (1.8).

We want to use an inductive argument to prove the thesis. If $\dim W = 0$ then this is Mukai's conjecture; it was proved by Peternell, Kollár, Ye-Zhang (see for instance [YZ]). Let the claim be true for dimension $m-1$. Note that the locus over which the fiber is not $\mathbf{P}^{r-1}$ is discrete and W has isolated singularities. In fact take a general hyperplane section A of W , and $X' = \pi^{-1}(A)$ then $\pi|_{X'} : X' \rightarrow A$ is again a contraction supported by $K_{X'} + \det E|_{X'}$, such that $r \geq ((n-1)+1)/2$. Thus by induction A is smooth, hence W has isolated singularities.

Let U be an open disk in the complex topology, such that $U \cap \text{Sing} W = \{0\}$. Then by lemma below 3.3 we have locally, in the complex topology, a π -ample line bundle L such that restricted to the general fiber is $\mathcal{O}(1)$. As in [Fu1, Prop. 2.12] we can prove that U is smooth and all the fibers are $\mathbf{P}^{r-1}$. $\square$

Lemma 3.3 *Let X be a complex manifold and $(W, 0)$ an analytic germ such that $W \setminus \{0\} \simeq \Delta^m \setminus \{0\}$. Assume we have an holomorphic map $\pi : X \rightarrow W$ with $-K_X$ π -ample; assume also that $F \simeq \mathbf{P}^r$ for all fibers of π , $F \neq F_0 = \pi^{-1}(0)$, and that $\text{codim} F_0 \geq 2$. Then there exists a line bundle L on X such that L is π -ample and $L|_F = \mathcal{O}(1)$.*

Proof. (see also [ABW2, pag 338, 339]) Let $W^* = W \setminus \{0\}$ and $X^* = X \setminus F_0$. By abuse of notation call $\pi = \pi|_{X^*} : X^* \rightarrow W^*$; it follows immediately that $R^1\pi_* \mathbf{Z}_{X^*} = 0$ and $R^2\pi_* \mathbf{Z}_{X^*} = \mathbf{Z}$.

If we look at Leray spectral sequence, we have that:

$$E_2^{0,2} = \mathbf{Z} \text{ and } E_2^{p,1} = 0 \text{ for any } p.$$

Therefore $d_2 : E_2^{0,2} \rightarrow E_2^{2,1}$ is the zero map and moreover we have the following exact sequence

$$0 \rightarrow E_\infty^{0,2} \rightarrow E_2^{0,2} \xrightarrow{d_3} E_2^{3,0},$$

since the only non zero map from $E_2^{0,2}$ is d_3 and hence $E_\infty^{0,2} = \ker d_3$. On the other hand we have also, in a natural way, a surjective map $H^2(X^*, \mathbf{Z}) \rightarrow E_\infty^{0,2} \rightarrow 0$. Thus we get the following exact sequence

$$H^2(X^*, \mathbf{Z}) \xrightarrow{\alpha} E_2^{0,2} \rightarrow E_2^{3,0} = H^3(W^*, \mathbf{Z}).$$

We want to show that α is surjective. If $\dim W := w \geq 3$ then $H^3(W^*, \mathbf{Z}) = 0$ and we have done. Suppose $w = 2$ then $H^3(W^*, \mathbf{Z}) = \mathbf{Z}$; note that the restriction of $-K_X$ gives a non zero class (in fact it is $r+1$ times the generator) in $E_2^{0,2}$ and is mapped to zero in $E_2^{3,0}$ thus the mapping $E_2^{0,2} \rightarrow E_2^{3,0}$ is the zero map and α is surjective. Since F_0 is of codimension at least 2 in X the restriction map $H^2(X, \mathbf{Z}) \rightarrow H^2(X^*, \mathbf{Z})$ is a bijection. By the vanishing of $R_i \pi_* \mathcal{O}_X$ we get $H^2(X, \mathcal{O}_X) = H^2(W, \mathcal{O}_W) = 0$ hence also $\text{Pic}(X) \rightarrow H^2(X, \mathbf{Z})$ is surjective. Let $L \in \text{Pic}(X)$ be a preimage of a generator of $E_2^{0,2}$. By construction L_t is $\mathcal{O}(1)$, for $t \in W^*$. Moreover $(r+1)L = -K_X$ on X^* thus, again by the codimension of X^* , this is true on X and L is π -ample.

□

4 An approach to the singular case

The following theorem arose during a discussion between us and J.A. Wisniewski; we would like to thank him. The idea to investigate this argument came from a preprint of Zhang [Zh2] where he proves the following result under the assumption that E is spanned by global sections. For the definition of log-terminal singularity we refer to [KMM].

Theorem 4.1 *Let X be an n -dimensional log-terminal projective variety and E an ample vector bundle of rank $n+1$, such that $c_1(E) = c_1(X)$. Then $(X, E) = (\mathbf{P}^n, \oplus^{n+1} \mathcal{O}_{\mathbf{P}^n}(1))$.*

Proof. We will prove that X is smooth, then we can apply proposition (3.2). We consider also in this case the associated projective space bundle Y and the commutative diagram

$$\begin{array}{ccc} \mathbf{P}(E) = Y & \xrightarrow{\varphi} & V \\ \downarrow p & & \downarrow \psi \\ X & \xrightarrow{\pi} & pt \end{array} \quad (3)$$

as in (1); it is immediate that Y is a weak Fano variety (i.e. Y is Gorenstein, log-terminal and $-K_Y$ is ample; in particular it has Cohen-Macaulay singularities); moreover, as in (3.1), $\dim F(\varphi) \leq \dim F(\pi) = n$ and the map φ is supported by $K_Y + (n+1)H$, where $H = \xi_E + A$, with ξ_E the tautological line bundle and A a pull back of an ample line bundle from V . It is known that a contraction supported by $K_Y + rH$ on a log terminal variety has to have fibers of dimension $\geq (r-1)$ and of dimension $\geq r$ in the birational case ([AW, remark 3.1.2]). Therefore in our case φ can not be birational and all fibers have dimension n ; moreover, by the Kobayashi-Ochiai criterion the general fiber is $F \simeq \mathbf{P}^n$. We want to adapt the proof of [BS, Prop 1.4]; to this end we have only to show that there are no fibers of φ entirely contained in $\text{Sing}(Y)$. Note that, by construction, $\text{Sing}(Y) \subset p^{-1}(\text{Sing} X)$ hence no fibers F of φ can be contained in $\text{Sing}(Y)$. Hence the same proof of [BS, Prop 1.4] applies and we can prove that V is nonsingular and $\varphi : Y \rightarrow V$ is a classical scroll. In particular Y is nonsingular and therefore also X is nonsingular. $\square$

More generally we can prove the following.

Theorem 4.2 *Let X be an n -dimensional log-terminal projective variety and E be an ample vector bundle of rank r . Assume that $K_X + \det E$ is nef and let $\pi : X \rightarrow W$ be the contraction supported by $K_X + \det E$. Assume also that for any fiber F of π $\dim F \leq r-1$, that $r \geq (n+1)/2$ and $\text{codim Sing}(X) > \dim W$. Then X is smooth and for any fiber $F \simeq \mathbf{P}^{r-1}$.*

Proof. The proof that X is smooth is as in the theorem above and then we use proposition (3.2) $\square$

5 Main theorem

This section is devoted to the proof of the following theorem.

Theorem 5.1 *Let X be a smooth projective variety over the complex field of dimension $n \geq 3$ and E an ample vector bundle on X of rank $r = (n-2)$. Then we have*

1) $K_X + \det(E)$ is nef unless (X, E) is one of the following:

i) there exist a smooth n -fold, W , and a morphism $\phi : X \rightarrow W$ expressing X as a blow up of a finite set B of points and an ample vector bundle E' on W such that $E = \phi^* E' \otimes [-\phi^{-1}(B)]$.

Assume from now on that (X, E) is not as in (i) above (that is eventually consider the new pair (W, E') coming from (i)).

- ii) $X = \mathbf{P}^n$ and $E = \oplus^{(n-2)} \mathcal{O}(1)$ or $\oplus^2 \mathcal{O}(2) \oplus^{(n-4)} \mathcal{O}(1)$ or $\mathcal{O}(2) \oplus^{(n-3)} \mathcal{O}(1)$ or $\mathcal{O}(3) \oplus^{(n-3)} \mathcal{O}(1)$.
- iii) $X = \mathbf{Q}^n$ and $E = \oplus^{(n-2)} \mathcal{O}(1)$ or $\mathcal{O}(2) \oplus^{(n-3)} \mathcal{O}(1)$ or $\mathbf{E}(2)$ with $\mathbf{E}$ a spinor bundle on $\mathbf{Q}^n$.
- iv) $X = \mathbf{P}^2 \times \mathbf{P}^2$ and $E = \oplus^2 \mathcal{O}(1, 1)$
- v) X is a del Pezzo manifold with $b_2 = 1$, i.e. $\text{Pic}(X)$ is generated by an ample line bundle $\mathcal{O}(1)$ such that $\mathcal{O}(n-1) = \mathcal{O}(-K_X)$ and $E = \oplus^{(n-1)} \mathcal{O}(1)$.
- vi) X is a classical scroll or a quadric bundle over a smooth curve Y .
- vii) X is a fibration over a smooth surface Y with all fibers isomorphic to $\mathbf{P}^{(n-2)}$.

2) If $K_X + \det(E)$ is nef then it is big unless there exists a morphism $\phi : X \rightarrow W$ onto a normal variety W supported by (a large multiple of) $K_X + \det(E)$ and $\dim(W) \leq 3$; let F be a general fiber of ϕ and $E' = E|_F$. We have the following according to $s = \dim W$:

- i) If $s = 0$ then X is a Fano manifold and $K_X + \det(E) = 0$. If $n \geq 6$ then $b_2(X) = 1$ except if $X = \mathbf{P}^3 \times \mathbf{P}^3$ and $E = \oplus^4 \mathcal{O}(1, 1)$.
- ii) If $s = 1$ then W is a smooth curve and ϕ is a flat (equidimensional) map. Then (F, E') is one of the pair described in [PSW], in particular F is either $\mathbf{P}^n$ or a quadric or a del Pezzo variety. If $n \geq 6$ then π is an elementary contraction. If the general fiber is $\mathbf{P}^{n-1}$ then X is a classical scroll while if the general fiber is $\mathbf{Q}^{n-1}$ then X is a quadric bundle.
- iii) If $s = 2$ and $n \geq 5$ then W is a smooth surface, ϕ is a flat map and (F, E') is one of the pair described in the Main Theorem of [Fu2]. If the general fiber is $\mathbf{P}^{n-2}$ all the fibers are $\mathbf{P}^{n-2}$.
- iv) If $s = 3$ and $n \geq 5$ then W is a smooth 3-fold and all fibers are isomorphic to $\mathbf{P}^{n-3}$.

3) Assume finally that $K_X + \det(E)$ is nef and big but not ample. Then a high multiple of $K_X + \det(E)$ defines a birational map, $\varphi : X \rightarrow X'$, which contracts an "extremal face" (see section 2). Let R_i , for i in a finite set of index, the extremal rays spanning this face; call $\rho_i : X \rightarrow W$ the contraction associated to one of the R_i . Then we have that each ρ_i is birational and divisorial; if D is one of the exceptional divisors (we drop the index) and $Z = \rho(D)$ we have that $\dim(Z) \leq 1$ and the following possibilities occur:

- i) $\dim Z = 0$, $D = \mathbf{P}^{(n-1)}$ and $D|_D = \mathcal{O}(-2)$ or $\mathcal{O}(-1)$; moreover, respectively, $E|_D = \oplus^{n-2}\mathcal{O}(1)$ or $E|_D = \oplus^{n-1}\mathcal{O}(1) \oplus \mathcal{O}(2)$.
- ii) $\dim Z = 0$, D is a (possible singular) quadric, $\mathbf{Q}^{(n-1)}$, and $D|_D = \mathcal{O}(-1)$; moreover $E|_D = \oplus^{n-2}\mathcal{O}(1)$.
- iii) $\dim Z = 1$, W and Z are smooth projective varieties and ρ is the blow-up of W along Z . Moreover $E|_F = \oplus^{n-2}\mathcal{O}(1)$.

If $n > 3$ then φ is a composition of "disjoint" extremal contractions as in i), ii) or iii).

Proof. Proof of part 1) of the theorem

Let (X, E) be a generalized polarized variety and assume that $K_X + \det(E)$ is not nef. Then there exist on X a finite number of extremal rays, $R_1, \dots, R_s$, such that $(K_X + \det(E)) \cdot R_i < 0$ and therefore, by the remark in section (2), $l(R_i) \geq (n-1)$.

Consider one of this extremal rays, $R = R_i$, and let $\rho : X \rightarrow Y$ be its associated elementary contraction. Then $L := -(K_X + \det(E))$ is ρ -ample and also the vector bundle $E_1 := E \oplus L$ is ρ -ample; moreover $K_X + \det(E_1) = \mathcal{O}_X$ relative to ρ . We can apply the theorem in [ABW2] which study the positivity of the adjoint bundle in the case of $\text{rank} E_1 = (n-1)$. More precisely we need a relative version of this theorem, i.e. we do not assume that E_1 is ample but that it is ρ -ample (or equivalently a local statement in a neighborhood of the exceptional locus of the extremal ray R). We just notice that the theorem in [ABW2] is true also in the relative case and can be proved exactly with the same proof using the relative minimal model theory (see [K-M-M]; see also the section 2 of the paper [AW] for a discussion of the local set up).

Assume first that ρ is birational, then $K_X + \det(E_1)$ is ρ -nef and ρ -big; note also that, since $l(R_i) \geq (n-1)$, ρ is divisorial. Therefore we are in the (relative) case C of the theorem in [ABW2] (see also the proposition 3.1 with $r = (n-1)$); this implies that Y is smooth and ρ is the blow up of a point in Y . Since $l(R_i) \geq (n-1)$, the exceptional loci of the birational rays are pairwise disjoint by proposition (1.6). This part give the point (i) of the theorem 5.1; i.e. the birational extremal rays have disjoint exceptional loci which are divisors isomorphic to $\mathbf{P}^{(n-1)}$ and which contract simultaneously to smooth distinct points on a n -fold W . The description of E follows trivially (see also [ABW2]).

If ρ is not birational then we are in the case B of the theorem in [ABW2]; from this we obtain similarly as above the other cases of the theorem 5.1, with some trivial computations needed to recover E from E_1 .

□

Proof of the part 2) of the theorem

Let $K_X + \det E$ be nef but not big; then it is the supporting divisor of a face $F = (K_X + \det E)^\perp$. In particular we can apply the theorems of section (2): therefore there exist a map $\pi : X \rightarrow W$ which is given by a high multiple of $K_X + \det E$ and which contracts the curves in the face. Since $K_X + \det E$ is not big we have that $\dim W < \dim X$. Moreover for every rational curve C in a general fiber of π we have $-K_X \cdot C \geq (n-2)$ by the remark in section (2). We apply proposition (1.8), which, together with the above inequality on $-K_X \cdot C$, says that π is an elementary contraction if $n \geq 5$ unless either $n = 6$, W is a point and X is a Fano manifold of pseudoindex 4 and $\rho(X) = 2$ or $n = 5$ and $\dim W \leq 1$.

By proposition (1.5) we have the inequality

$$n + \dim F \geq n + n - 2 - 1;$$

in particular it follows that $\dim W \leq 3$.

(5.1) Let $\dim W = 0$, that is $K_X + \det E = 0$ and therefore X is a Fano manifold. By what just said above we have that $b_2(X) = 1$ if $n \geq 6$ with an exception which will be treated in the following lemma.

Lemma 5.2 *Let X be a 6 dimensional projective manifold, E is an ample vector bundle on X of rank 4 such that $K_X + \det E = 0$. Assume moreover that $b_2 \geq 2$. Then $X = \mathbf{P}^3 \times \mathbf{P}^3$ and $E = \oplus^4 \mathcal{O}(1, 1)$.*

Proof. The lemma is a slight generalization of [Wil, Prop B] for dimension 6; the poof is similar and we refer to this paper. In particular as in [Wil] we can see that X has two extremal rays whose contractions, $\pi_i, i = 1, 2$, are of fiber type with equidimensional fibers onto 3-folds W_i and with general fiber $F_i \simeq \mathbf{P}^3$. We claim that the W_i are smooth and thus $W_i \simeq \mathbf{P}^3$. First of all note that W_i can have only isolated singularity and only isolated points over which the fiber is not $\mathbf{P}^{n-3}$; in fact let S be a general hyperplane section of W_i and $T_i = \pi_i^*(S)$, then $(\pi_i)_{|T_i}$ is an extremal contraction, by proposition 1.8; hence by [ABW2, Prop 1.4.1] S is smooth; moreover the contraction is supported by $K_{T_i} + \det E_{T_i}$ hence all fibers are $\mathbf{P}^3$ by the main theorem of [ABW2]. Now we are (locally) in the hypothesis of lemma 3.3 so we get, locally in the complex topology, a tautological bundle and we can conclude, by [Fu1, Prop 2.12], that W_i is smooth. Let $T = H_1 \cap H_2$, where H_i are two general elements of $\pi_1^*(\mathcal{O}(1))$. T is smooth, we claim that $T \simeq \mathbf{P}^1 \times \mathbf{P}^3$. In fact $\pi_{1|T}$ makes T a projective bundle over a line (since $H^2(\mathbf{P}^1, \mathcal{O}^*) = 0$), that is $T = \mathbf{P}(\mathcal{F})$. Moreover $\pi_{2|T}$ is onto $\mathbf{P}^3$, therefore the claim follows. Therefore we conclude that $\pi_i^* \mathcal{O}_{\mathbf{P}^3}(1)_{|F_i} \simeq \mathcal{O}_{\mathbf{P}^3}(1)$ for $i = 1, 2$. This implies by Grauert Theorem that the two fibrations are classical scroll, that is $X = \mathbf{P}(\mathcal{F}_i)$, for $i = 1, 2$; moreover computing the canonical class of X the $\mathcal{F}_i$ are ample and the lemma easily follows.

□

(5.2) Let $\dim W = 1$. Then W is a smooth curve and π is a flat map. Let F be a general fiber, then F is a smooth Fano manifold and $E|_F$ is an ample vector bundle on F of rank $(n-2) = \dim F - 1$ such that $-K_F = \det(E|_F)$. These pairs $(F, E|_F)$ are classified in the Main Theorem of [PSW]; in particular if $\dim F \geq 5$ F is either $\mathbf{P}^{(n-1)}$ or $\mathbf{Q}^{(n-1)}$ or a del Pezzo manifold with $b_2(F) = 1$. Moreover if $n \geq 6$ then π is an elementary contraction by proposition (1.8).

Claim Let $n \geq 6$ and assume that the general fiber is $\mathbf{P}^{n-1}$, then X is a classical scroll and $E|_F$ is the same for all F .

(See also [Fu2]) Let $S = W \setminus U$ be the locus of points over which the fiber is not $\mathbf{P}^{n-1}$. Over U we have a projective fiber bundle. Since $H^2(U, \mathcal{O}^*) = 0$ we can associate this $\mathbf{P}$ -bundle to a vector bundle $\mathcal{F}$ over U . Let $Y = \mathbf{P}(\mathcal{F})$ and H the tautological bundle; by abuse of language let H the extension of H to X . Since π is elementary H is an ample line bundle on X . Therefore by semicontinuity $\Delta(F, H_F) \geq \Delta(G, H_G)$, for any fiber G , where $\Delta(X, L)$ is Fujita delta-genus. In our case this yields $0 = \Delta(F, H_F) \geq \Delta(G, H_G) \geq 0$. Moreover by flatness $(H_G)^{n-1} = (H_F)^{n-1} = 1$ and Fujita classification allows to conclude. The possible vector bundle restricted to the fibers are all decomposables, hence they are rigid, that is $H^1(\text{End}(E)) = \oplus_i H^1(\text{End}(\mathcal{O}(a_i))) = \oplus_i H^1(\mathcal{O}(-a_i)) = 0$. Hence the decomposition is the same along all fibers of π .

Claim Let $n \geq 6$ and assume that the general fiber is $\mathbf{Q}^{n-1}$. Then X is a quadric bundle.

Let as above $S = W \setminus U$ be the locus of points over which the fiber is not a smooth quadric. Let $X^* = \pi^{-1}(U)$ then we can embed X^* in a fiber bundle of projective spaces over U , since it is locally trivial. Associate this P -bundle over U to a projective bundle and argue as before.

□

(5.3) Let now $\dim W = 2$ and assume that $n \geq 5$; then π is an elementary contraction. This implies first, by [ABW2, Prop. 1.4.1], that W is smooth; secondly that π is equidimensional, hence flat and the general fiber is $\mathbf{P}^{n-2}$ or $\mathbf{Q}^{n-2}$, see [Fu2].

Claim Let $n \geq 5$ and the general fiber is $\mathbf{P}^{n-2}$ then for any fiber $F \simeq \mathbf{P}^{n-2}$ and $E|_F$ is the same for all F .

Let $S \subset W$ be the locus of singular fibers, then $\dim S \leq 0$ since W is normal. Let $U \subset W$ be an open set, in the complex topology, with $U \cap S = \{0\}$ and let $V \subset X$ such that $V = \pi^{-1}(U)$. We are in the hypothesis of lemma 3.3 thus we get a "tautological" line bundle H on V and we conclude by [Fu1, Prop. 2.12].

There are two possible restriction of E to the fiber, namely $E|_F \simeq \mathcal{O}(2) \oplus (\oplus^{n-1} \mathcal{O}(1))$ or $E|_F$ is the tangent bundle. As observed by Fujita in [Fu2] this two restrictions have a different behavior in the diagram (1), in the former φ is

birational while in the latter it is of fiber type. Hence the restriction has to be constant along all the fibers. $\square$

(5.4) Let finally $\dim W = 3$; the general fiber is $\mathbf{P}^{n-3}$ (see for instance [Fu2]). Assume that $n \geq 5$, therefore π is elementary; we claim that all fibers are $\mathbf{P}^{n-3}$.

Since π is elementary any fiber G has $\text{cod}G \geq 2$. Let $S \subset W$ be the locus of point over which the fiber is not $\mathbf{P}^{n-3}$; $\dim S \leq 0$ since a generic linear space section can not intersect S , by the above. Let $U \subset W$ be an open set, in the complex topology, with $U \cap S = \{0\}$ and let $V \subset X$ such that $\pi(V) = U$. Then by lemma 3.3 we get a "tautological" line bundle H on V ; $\pi : V \rightarrow U$ is supported by $K_V + (n-2)H$. Thus by [AW, Th 4.1] U is smooth and all the fibers are $\mathbf{P}^{n-3}$ (we use that $n \geq 5$). $\square$

Proof of the part 3) of the theorem

In the last part of the theorem we assume that $K_X + \det E$ is nef and big but not ample. Then $K_X + \det E$ is a supporting divisor of an extremal face, F ; let R_i the extremal rays spanning this face. Fix one of this ray, say $R = R_i$ and let $\pi : X \rightarrow W$ be the elementary contraction associated to R .

We have $l(R) \geq n-2$; this implies first that the exceptional loci are disjoint if $n > 3$, proposition (1.7). Secondly, by the inequality (1.5), we have

$$\dim E(R) + \dim F(R) \geq 2n - 3.$$

Therefore $\dim E(R) = n-1$ and either $\dim F(R) = n-1$ or $\dim F(R) = n-2$; if $Z := \rho(E)$ and $D = E(R)$ this implies that either $\dim Z = 0$ or 1.

If $\dim Z = 1$ then $\dim F(\pi) = n-2$ for all fibers (note that since the contraction π is elementary there cannot be fiber of dimension $(n-1)$); thus we can apply proposition (3.1) with $r = (n-2)$. This will give the case 3-(iii) of the theorem.

Consider again the construction in section (2), in particular we refer to the diagram (1). Let S be the extremal ray contracted by φ ; note that $l(S) \geq n-2$ and that the inequality (1.5) gives

$$\dim E(S) + \dim F(S) \geq 3n - 6;$$

in particular, since $\dim F(S) \leq \dim F(R)$, we have two cases, namely $\dim E(S) = 2n-5$ and $\dim F(S) = (n-1)$ or $\dim E(S) = 2n-4$ and $\dim F(S) = (n-1)$ or $(n-2)$.

The case in which $\dim E(S) = 2n-5$ will not occur. In fact, after "slicing", (see 2), we would obtain a map $\varphi' = \varphi|_Z$ which would be a small contraction supported by a divisor of the type $K_Z + (n-2)L$ but this is impossible by the classification of [Fu1, Th 4] (see also [An]).

Hence $\dim E(S) = 2n - 4$, that is also φ is divisorial.

Suppose that the general fiber of φ , $F(S)$, has dimension $(n - 2)$. After slicing we obtain a map $\varphi' = \varphi|_Z : Z \rightarrow T$ supported by $K_Z + (n - 2)L$, where $L = \xi_{E|Z}$. This map contracts divisors D in Z to curves; by ([Fu1, Th 4]) we know that every fiber F of this map is $\mathbf{P}^{(n-2)}$ and that $D|_F = \mathcal{O}(-1)$ (actually this map is a blow up of a smooth curve in a smooth variety). In particular there are curves in Y , call them C , such that $-E(S) \cdot C = 1$. We will discuss this case in a while.

Suppose then the general fiber of φ , $F(S)$, has dimension $(n - 1)$; therefore all fibers have dimension $(n - 1)$. Slicing we obtain a map $\varphi' = \varphi|_Z : Z \rightarrow T$ supported by $K_Z + (n - 2)L$, where $L = \xi_{E|Z}$. This map contracts divisors D in Z to points; by ([Fu1]) we know that these divisors are either $\mathbf{P}^{(n-1)}$ with normal bundle $\mathcal{O}(-2)$ or $\mathbf{Q}^{(n-1)} \subset \mathbf{P}^n$ with normal bundle $\mathcal{O}(-1)$. In the latter case we have as above that there are curves C in Y , such that $-E(S) \cdot C = 1$.

In these cases observe that $E(S) \cdot \tilde{C} = 0$, where $\tilde{C}$ is a curve in the fiber of p . Hence $E(S) = p^*(-M)$ for some $M \in \text{Div}(X)$. Let l be an extremal curve of $E(S)$. Then, by projection formula, we have $-1 = E(S) \cdot l = -M \cdot mC$ and thus M generates $\text{Im}[Pic(X) \rightarrow Pic(D)]$, hence M is π -ample; note that in general it does not generate $Pic(D)$. We study now the Hilbert polynomial of $M|_D$ to show that $\Delta(D, M|_D) = 0$, where $\Delta(X, L)$ is Fujita delta genus. Let $\mathcal{O}_D(-K_X) \simeq \mathcal{O}_D(pM)$, where $p = l(R) \geq n - 2$, and $\mathcal{O}_D(-D) \simeq \mathcal{O}_D(qM)$ for some $p, q \in \mathbf{N}$. By adjunction formula $\omega_D \simeq \mathcal{O}_D(-(p + q)M)$. By [Ando, Lemma 2.2] or [BS, pag 179], Serre duality and relative vanishing we obtain that $q \leq 2$, the Hilbert polynomial is

$$P(D, M|_D) = \frac{a}{(n-1)!} (t+1) \cdots (t+(n-2))(t+c)$$

and the only possibilities are $a = 1, c = n - 1, q = 1 \text{ or } 2$ and $a = 2, c = (n - 1)/2, q = 1$. In particular $\Delta(D, M|_D) = 0$ and, by Fujita classification, D is equal to $\mathbf{P}^{(n-1)}$ or to $\mathbf{Q}^{(n-1)} \subset \mathbf{P}^n$. Now the rest of the claim in 3) i) and ii) follows easily.

It remains the case in which $\varphi' = \varphi|_Z : Z \rightarrow T$ contracts divisors $D = \mathbf{P}^{(n-1)}$ with normal bundle $\mathcal{O}(-2)$ to points. We can apply the above proposition (2.1) and show that the singularities of W are the same as those of T . Then, as in ([Mo1]), this means that we can factorize π with the blow up of the singular point. Let $X' = Bl_w(W)$, then we have a birational map $g : X \rightarrow X'$. Note that X' is smooth and that g is finite. Actually it is an isomorphism outside D and cannot contract any curve of D . Assume to the contrary that g contracts a curve $B \subset D$; let $N \in Pic(X')$ be an ample divisor then we have $g^*N \cdot B = 0$

while $g^*N \cdot C \neq 0$ contradiction. Thus by Zarisky's main theorem g is an isomorphism. This gives a case in 3)i).

References

- [Ando] Ando, T., On extremal rays of the higher dimensional varieties, *Inv.Math.* **81** (1985), 347-357
- [An] Andreatta,M., Contractions of Gorenstein polarized varieties with high nef value, to appear on *Math. Ann.* (1994).
- [ABW1] Andreatta,M.- Ballico,E.-Wiśniewski, On contractions of smooth algebraic varieties, preprint UTM 344 (1991).
- [ABW2] Andreatta,M.- Ballico,E.-Wiśniewski, Vector bundles and adjunction, *International Journal of Mathematics*,**3** (1992), 331-340.
- [AW] Andreatta,M.- Wiśniewski, J., A note on non vanishing and its applications,*Duke Math.J.*, **72**, (1993).
- [BS] Beltrametti,M. Sommese,A.J. On the adjunction theoretic classification of polarized varieties, *J. reine angew. Math* **427** (1992), 157-192.
- [CKM] Clemens,H. Kollár,J. Mori,S. Higher dimensional complex geometry *Asterisque* **166** 1988
- [Fu1] Fujita, T., On polarized manifolds whose adjoint bundle is not semipositive, in *Algebraic Geometry, Sendai*, *Adv. Studies in Pure Math.* **10**, Kinokuniya–North-Holland 1987, 167—178.
- [Fu2] Fujita, T., On adjoint bundle of ample vector bundles, in *Proc. Alg. Geom. Conf. Bayreuth* (1990), *Lect. Notes Math.*, **1507**, 105-112.
- [Fu3] Fujita,T. On Kodaira energy and reduction of polarized manifolds, *Manuscr. Math* **76** (1992), 59-84.
- [KMM] Kawamata, Y., Matsuda, K., Matsuki, K., Introduction to the Minimal Model Program in Algebraic Geometry, Sendai, *Adv. Studies in Pure Math.* **10**, Kinokuniya–North-Holland 1987, 283-360.
- [Io] Ionescu,P. Generalized adjunction and applications, *Math. Proc. Camb. Phil. Soc.* **99** (1986), 457-472.
- [Mo] Mori,S. Projective manifolds with ample tangent bundle, *Ann.Math.* **110** (1979), 593-606
- [Mo1] Mori,S., Threefolds whose canonical bundles are not numerical effective, *Ann. Math.*, **116**, (1982), 133-176.
- [Mu] Mukai,S. Problems on characterizations of complex projective space, *Birational Geometry of Algebraic varieties - Open problems*, Katata Japan (1988), 57-60.
- [OSS] Okonek,C. Schneider,M. Spindler,H. Complex vector bundle on projective space. (*Prog.Math*) Boston Basel Stuttgart: Birkhäuser 1981

- [PSW] Peternell, T.- Szurek, M.- Wiśniewski, J.A., Fano Manifolds and Vector Bundles, *Math. Ann.*, **294**, (1992), 151-165.
- [So] Sommese, A.J., On the adjunction theoretic structure of projective varieties, *Complex Analysis and Algebraic Geometry*, Proceedings Göttingen, 1985 (ed. H. Grauert), *Lecture Notes in Math.*, **1194** (1986), 175—213.
- [Wi1] Wiśniewski, J.A., On a conjecture of Mukai, *Manuscr. Math.* **68** (1990), 135-141.
- [Wi2] Wiśniewski, J.A., Length of extremal ray and generalized adjunction, *Math. Z.* **200** (1989), 409-427.
- [Wi3] Wiśniewski, J.A., On contraction of extremal rays of Fano manifolds, *J. reine und angew. Math.* **417** (1991), p. 141-157.
- [YZ] Ye, Y.G. - Zhang, Q., On ample vector bundle whose adjunction bundles are not numerically effective, *Duke Math. Journal*, **60** n. 3 (1990), p. 671-687.
- [Zh] Zhang, Q., A theorem of the adjoint system for vector bundles, *Manuscripta Math.*, **70** (1991), p. 189-201.
- [Zh2] Zhang, Q., Ample vector bundle, preprint